

*If there is a chance you could be pregnant, please tell the tech.*

## Renal Scan With Furosemide (Lasix)

A renal scan is a test that shows the size, shape and function of your kidneys. If your kidney function is decreased, a second step is added to the renal scan to check how your kidneys respond to a diuretic medication or “water pill.” Furosemide (Lasix®) is the diuretic medication given to increase the amount of urine your kidneys make. The amount of radiation used in this test is small and well within limits that are not harmful to you. This test is done in the Nuclear Medicine Department and lasts about 2 1/2 hours.

### Before the test

There is no preparation you need to do for this test.

On the day of the test, plan to arrive 15 minutes before the test is scheduled to begin. Be sure to bring:

- Your physician’s written order for the test if you received one
- A list of allergies
- A list of all your current medications (prescription, over-the-counter and herbal)
- Photo ID
- Medical insurance information
- Medicare card (Medicare patients only)

Please come to the 8th floor Reception Desk in Galter Pavilion, 201 East Huron Street.

Parking is available for patients and visitors in the garage at 222 East Huron Street, across from Feinberg and Galter pavilions. For a discounted rate, please bring your parking ticket with you. Your ticket can be validated at the Customer Services Desks on the 1st, 2nd and 8th floors of Feinberg and Galter pavilions, and the 1st floor of Prentice Women’s Hospital.

### During the test

When you arrive in the Nuclear Medicine Department, a nurse or technologist (tech) may ask you questions about your medical history. *If you think you might be pregnant or are breastfeeding, please tell the tech before the start of the test and before you are given an injection.*

The tech will ask you to lie on an imaging table. A radioactive tracer will be injected by an IV (intravenous) line into a vein in your arm or hand. A special camera will detect the flow of the tracer to your kidneys. Pictures of your kidneys and bladder will be taken for 30 minutes. At the end of this time, the tech will ask you to empty your bladder, and another picture of your kidneys will be taken.

The tech will check your pictures with a physician. Then, a physician, nurse or tech will give you Lasix through your IV line. Pictures of your kidneys and bladder are repeated for another 30 minutes. The tech will then ask you to empty your bladder again.

A last picture of your kidneys and bladder will be taken as you stand in front of the camera.

### **After the test**

You may resume your normal activities and diet after the test. The radioactive tracer you were given is not harmful and becomes nonradioactive within hours. The Lasix injection may cause you to urinate frequently for a couple of hours after the test.

Talk with your physician about your test results. You may obtain a copy of your test results about a week after the test from the Medical Records Department, which is located on the 2nd floor of Galter Pavilion.

If you have any questions about this test, please call the Nuclear Medicine Department at 312.926.2320, TTY 312.926.6363, Monday through Friday, 7:30 am to 5:00 pm.