

## Cardiac Surgery: Heart Valve Disease


*If you have  
questions, ask  
your care team.*

Your care team may have talked about the need for heart valve surgery with you. This brochure will help you better understand heart valve surgery and what to expect. It will explain how the heart valves work, the types of valve disease and the surgeries to treat them.


### Heart valves

To understand heart valve disease, it is helpful to know how the valves in the heart work. There are 4 heart valves: mitral, tricuspid, aortic and pulmonary. Valves have either 2 or 3 leaflets (flaps). Each valve opens and closes about 50 million times a year, up to 4 billion times in a typical lifetime. The valves keep the blood moving forward through the heart and out to the rest of the body. In a front view of the heart, Figures 1 and 2 show normal-looking mitral and aortic valves. These are the valves most often affected by disease.

**Figure 1. Closed mitral valve**



**Figure 2. Closed aortic valve**



©2015 Krames Staywell LLC

When the mitral valve opens on the left side of the heart, it lets blood into the left ventricle (lower heart chamber). The aortic valve is closed. The left ventricle then fills with blood (Figure 2). When the mitral valve closes, the aortic valve opens and blood is pumped out to the rest of the body through the aorta (Figure 1).

Likewise, when the tricuspid valve opens on the right side of the heart, the right ventricle fills with blood. When this valve closes, the pulmonary valve opens and blood is pumped into the blood vessels of the lungs. This let the blood fill with oxygen before continuing its journey to the left ventricle and out to the rest of the body (Figure 3).

**Figure 3. Normal heart valves**




Figure 3 shows normally functioning valves as viewed from above the heart. Normally, 1 set of valves opens while the other set of valves closes.

In the top view, the pulmonary and aortic valves are closed. The mitral and tricuspid valves are wide open. This allows blood to fill the ventricles.

The lower picture shows the pulmonary and aortic valves completely open. The mitral and tricuspid valves are now securely closed. At this time, blood is pumped out of the ventricles.

### Heart valve disease

In heart valve disease, the valve leaflets do not open or close properly. This affects the blood flow. Valve regurgitation (insufficiency or leaking) and valve stenosis (narrowing or obstruction) are 2 common heart valve problems. Both regurgitation and stenosis cause the heart to work harder. This may lead to heart failure. Each of these conditions is briefly explained below.

#### Valve regurgitation


Regurgitation happens when the valve does not close tightly. This causes blood to flow backward instead of forward.

The 3 small yellow arrows in Figure 4 show the back flow of blood through a diseased mitral valve.

This happens at the same time the heart is trying to pump the blood out into the body (large yellow arrow).

\*©2010 Elsevier

**Figure 4. Back flow of blood**



©2013 Abbott Laboratories. All Rights Reserved.


### ***Valve stenosis***

Stenosis happens when the valve does not fully open. This limits how much blood flows through. Figure 5 shows how the valve leaflets change in appearance.

The top pictures show a normally functioning aortic valve. On the top left side, the valve is wide open to let the blood flow out. On the top right side, the valve is closed when it is at rest.

Below that, the aortic valve cannot fully open and close. This makes it harder for the blood to pass through, causing it to back up.

**Figure 5. Normal and stenotic aortic valve**



\*\* ©2007 Edwards Lifesciences LLC

### ***Causes of valve disease***

Valve disease may have many different causes, such as:

- Degenerative disease
- Calcium buildup on and around the valve leaflets
- Bicuspid aortic disease (present at birth)
- Genetic disorders (born with valve disease)
- Heart damage
- Cardiomyopathy (enlarged heart)
- Rheumatic heart disease
- Endocarditis (infection)
- Heart tumors

Your physician will discuss your specific cause with you to help you understand your treatment options.

### ***Symptoms of valve disease***

Symptoms of valve disease may occur suddenly or develop gradually. They may include:

- Fatigue
- Chest pain
- Palpitations
- Feeling dizzy or faint
- Swelling in the ankles and feet
- Shortness of breath (may get worse during activity or when lying down)

For more information about valve disease and the Northwestern Medicine Center for Heart Valve Disease, please go to [heartvalvedisease.nm.org](http://heartvalvedisease.nm.org).

### **Assessing patients with valve disease**

Before planning surgery, you will need to come to the Center for Heart Valve Disease. Expert physicians and nurses will work with you. They will talk with you in detail about your medical history, your valve disease and any symptoms you may be having. You will also have an in-depth physical exam that focuses on heart murmurs and other signs of valve stenosis or regurgitation.

Your physician may also want to see other test results that show how well your heart and valves function. These include:

- Echocardiogram (echo). An echo is an ultrasound of the heart. It is the most common test used to evaluate heart valve disease. It shows real-time heart and valve function. People often have an echo before and after an exercise test to show how the heart responds to stress.
- Transesophageal echo (TEE). In some cases, you may need this test for an even clearer look at your heart valves.
- Computed tomography (CT) scan. This test shows a detailed 3D image of the heart.
- Magnetic resonance imaging (MRI) scan. The MRI shows a more detailed look at how well the heart chambers work and how much valve regurgitation exists. Sometimes the physician will use an MRI instead of a CT scan to check valve disease.
- Cardiac catheterization. This test precisely defines any blockages in the coronary arteries. It also measures the pressures in the heart. For this test, the physician inserts a small catheter into the wrist or groin and threads it up to the heart.

For more information on testing, please refer to the “Pre-surgery evaluation” section of the Northwestern Memorial Hospital brochure *Before Heart Surgery: A Patient Guide*.


## Surgical treatment of valve disease

Treatment depends on the symptoms and severity of the disease. Treatment may include medication or surgery. Surgery may be either a valve repair or a valve replacement.


### Valve repair

The heart surgeons will repair a valve whenever possible. This often gives the best long-term outcome in restoring proper valve function. During heart valve repair, the surgeon trims, reshapes or rebuilds your valve. The care team places a ring at the base of the heart valve to for more support to the repaired valve (Figure 7). This procedure is known as an annuloplasty. Figures 6 and 7 show the mitral valve before and after it is repaired. You will need to take an anticoagulant medication for at least 3 months **after** valve repair surgery.

**Figure 6. Diseased valve**



**Figure 7. Valve after annuloplasty**



### Valve replacement

If the valve cannot be repaired, you will need a new valve. Your surgeon will remove the diseased valve and sew a new one into place. There are 2 main types of replacement valves: bioprosthetic (tissue) valves or mechanical valves. Valves come in different sizes and materials.

**Bioprosthetic (tissue) valve:** Tissue valves are made from animal or human tissue. The durability of a valve depends on your age at the time of implant. If you get the valve at a young age, it is more likely that you will need another valve in the future. Figure 8 shows one type of tissue valve.

You will need to take anticoagulant medication for 3 months **after** tissue valve surgery.

**Figure 8. Bioprosthetic valve**



\*\*©2007 Edwards Lifesciences LLC

**Mechanical valve:** Mechanical valves (Figure 9) are designed to last a lifetime. However, there is a risk of blood clots with these valves, which increases with each year after surgery. Blood clots may lead to strokes or other serious health problems. Therefore, anticoagulation (“blood-thinning”) medications are needed indefinitely if you get this type of valve. This may require some changes to your diet and to any physical activity, if you are active. If you become unable to take these medications, you would need surgery to replace the mechanical valve with a tissue valve.

**Figure 9. Mechanical valve**



©2007 On-X Life Technologies Inc.

### ***Alternatives to open heart surgery***


The valve surgeries above are considered “open heart surgeries.” This is because the surgeon makes an incision either through the chest or rib cage to directly access the heart. This is the conventional method.

If open heart surgery is not a good option for you, there are alternatives. In this case, your physician may recommend one of the following procedures.

**Transcatheter aortic valve replacement (TAVR):** In a TAVR procedure, the cardiologist replaces the diseased aortic valve with a bioprosthetic valve. Then, they insert a tube (catheter) through an artery in the groin or through a small incision between the ribs. The catheter is threaded up the artery until it reaches the heart. Then, they insert the new valve within the old valve. Figure 10 shows the new valve in place after a TAVR procedure.


More information about TAVR is available at [heartvalvedisease.nm.org/transcatheter-aortic-valve-replacement.html](http://heartvalvedisease.nm.org/transcatheter-aortic-valve-replacement.html).

**Figure 10. Implanted transcatheter aortic valve**



**Transcatheter mitral valve repair:** In the transcatheter mitral valve repair (MitraClip®), the cardiologist uses a “clip” to join the mitral valve leaflets together. (Figure 11). This reduces the backward flow of blood and lets the heart work more efficiently.

**Figure 11. MitraClip procedure**



©2013 Abbott Laboratories. All Rights Reserved.

For this procedure, the cardiologist inserts a catheter through a vein in the groin. The catheter is threaded up to the heart. Figure 11 shows how the catheter is passed through the septum of the heart to insert the clip. Ultrasound imaging confirms the placement. The cardiologist may place multiple clips, if needed. More information about the MitraClip procedure is available at [heartvalvedisease.nm.org/mitraclip-for-mitral-regurgitation.html](http://heartvalvedisease.nm.org/mitraclip-for-mitral-regurgitation.html).

### ***Valve surgery risks***

Every surgery carries some risk. The amount depends on factors such as your age and overall health. Risks may include bleeding, infection, and lung or heart problems. Atrial fibrillation (a type of irregular heart rhythm) is an example of a heart problem that may happen after surgery. In some cases, a pacemaker or other procedures may be needed. In rare instances, stroke or kidney failure may occur. Your cardiologist will discuss your risks with you.

### **After surgery: follow-up care**

#### **Follow-up care is required after all valve repair and valve replacements.**

Please read the Northwestern Memorial Hospital home care brochure, *Heart Surgery: Care After Leaving the Hospital*, to understand all parts of your follow-up care. Below are a few key points.

#### ***Blood-thinner medications and weekly blood tests***

If you are taking blood thinners (anticoagulant medication) you will need weekly blood tests after you go home. You will have these tests until your physician no longer needs to adjust your dose. Please refer to the medication brochure that your care team will give you, which has important information about blood-thinner medication.

### ***Dental/surgical procedures***

To help prevent infection to your heart valve, follow these guidelines:

- You should not have any dental procedures for 12 weeks after valve surgery.
- You will always need to take antibiotics before dental or surgical procedures in the future.
- Inform all clinicians and dental staff that you have had heart valve surgery, before you need any dental or surgical procedures.
- Tell your dentist that your heart surgeon suggests following the *American College of Cardiology/American Heart Association Valvular Heart Disease Guidelines*.

**Appendix A in the *Heart Surgery: Care After Leaving the Hospital* brochure will give you more information about long-term follow-up care after heart valve surgery.**

### **Bicuspid aortic valve**

If you have bicuspid aortic valve (BAV), please read the following. It is important to understand this disease and how it may affect you and your family.


The aortic valve controls the flow of blood from the heart out to the body. Normally, the aortic valve has 3 leaflets that keep blood moving in one direction. With BAV, 2 of the 3 leaflets of the aortic valve fuse together before birth. This creates a 2-leaflet valve, instead of the normal 3-leaflet valve (Figure 12).

BAV also affects the thoracic aorta, the largest artery in the body that carries blood from the heart to the rest of the body.

A heart with BAV may work well without causing problems for a long time. However, many people will eventually develop complications. Then, they will need surgery either to fix the aortic valve, the thoracic aorta or both.

BAV is one of the most common congenital heart defects present at birth. It is at least 2 times as likely to happen in men as in women. BAV may be inherited in families.

**Figure 12. Tricuspid and bicuspid aortic valves**



©alila/www.fotosearch.com k7918328

### ***Family screening for BAV***

If you have BAV, your family members may have BAV too. This makes family screening very important because most people with BAV have no symptoms until they start to have complications. Close family members (parents, siblings, children) should have an echo to see if they have BAV. An echo is a test that uses ultrasound waves to get real-time images of the heart and heart valve function.

For more information about the Northwestern Medicine BAV Program, go to **[bav.nm.org](http://bav.nm.org)**.

For more information about family screening or to make an appointment, please contact the Northwestern Medicine BAV nurse coordinator, at [bav\\_rn@nm.org](mailto:bav_rn@nm.org) or call 312.926-7410 (TTY: 711).

\*Carpentier's Reconstructive Valve Surgery, p. 27, Alain Carpentier, MD, PhD, David Adams, MD and Farzan Filsoufi, MD, Saunders (2010) Elsevier, Philadelphia, PA. Permission for use obtained.

\*\*Edwards, Edwards Lifesciences, Edwards SAPIEN, SAPIEN, SAPIEN XT, SAPIEN 3 are trademarks of Edwards Lifesciences Corporation, Irvine, CA. 2007. Permission for use obtained.